

ON A PAPER OF Ž. TOMOVSKI AND R. HILFER

VIKTOR ZASTAVNYI

ABSTRACT. The proof of the inequalities for alternating Mathieu type series in [2] by Ž. Tomovski and R. Hilfer contains a mistake. Here we give the values of parameters for which these inequalities are not true.

In paper [2] stated that under validity one of the conditions

- (1) **1)** $\beta = 1, \alpha = 2, \mu = 2$; **2)** $\beta = 0, \alpha = 2, \mu > \frac{1}{2}$; **3)** $\beta = 1, \alpha = 2, \mu > 1$;
4) $\beta > 0, \alpha \in \mathbb{N}, \mu\alpha - \beta > 1$; **5)** $\beta > 0, \alpha > 0, \mu\alpha - \beta > 1$,

the next inequality (see in [2] inequalities (2.17), (2.22), (2.23), (2.25), (2.27))

$$(2) \quad \sum_{k=1}^{\infty} \frac{2(-1)^{k-1}k^{\beta}}{(k^{\alpha} + r^2)^{\mu}} \leq \frac{2}{(1 + r^2)^{\mu}} , \quad r > 0 ,$$

take place.

The proof is based on the inequality

$$(3) \quad \int_0^{\infty} \frac{x^{\mu\alpha-\beta-1}}{e^x + 1} K(rx^{\alpha/2}) dx \leq \int_0^{\infty} x^{\mu\alpha-\beta-1} e^{-x} K(rx^{\alpha/2}) dx , \quad r > 0 ,$$

in the each of five cases. The last inequality (3) is not valid in general case, because the kernel $K(u)$ (which depends on the parameters β, α and μ) is not positive. For example, in the cases **1)**, **2)** and **3)** the kernel $K(u)$ is equal $\frac{\sin u}{u}$, $c_1(\mu)j_{\mu-\frac{1}{2}}(u)$, $c_2(\mu)j_{\mu-\frac{3}{2}}(u)$ correspondingly, where $c_1(\mu) > 0$, $c_2(\mu) > 0$, $j_{\lambda}(u) = \frac{J_{\lambda}(u)}{u^{\lambda}}$, J_{λ} - Bessel function of the first kind. In the cases **4)** and **5)** the positiveness of the kernel $K(u)$ is not obvious and rather it is alternating sign (at least for the parameters in the counterexample below). Consequently, in the five cases, the inequality (2) is not proved. We have to mention that for $\beta = 0, \alpha > 0, \mu > 0$ the strict inequality (2) is obvious.

Let us consider the values of the parameters for which the inequality (2) is not valid. In 2008, the author proved (see [3, Theorem 5] or [4, Theorem 2.4]), that if $(\gamma, \alpha) \in \mathbb{Z}_+ \times \mathbb{N}$ and $\alpha(\mu+1) - \gamma > 0$, then for every fixed $u \in \mathbb{R}$ the next asymptotic representation

$$\sum_{k=1}^{\infty} \frac{2(-1)^{k-1}(k+u)^{\gamma}}{((k+u)^{\alpha} + t^{\alpha})^{\mu+1}} \sim \sum_{k=0}^{\infty} \frac{(-1)^{k(\alpha+1)+\gamma}}{t^{\alpha(k+\mu+1)}} \cdot \frac{\Gamma(\mu+k+1) E_{k\alpha+\gamma}(-u)}{\Gamma(\mu+1)\Gamma(k+1)} , \quad t \rightarrow +\infty$$

is valid. Here $E_n(x)$ - Euler polynomials and $E_{n-1}(x) = \frac{2}{n} (B_n(x) - 2^n B_n(\frac{x}{2}))$, $n \in \mathbb{N}$, where $B_n(x)$ - Bernoulli polynomials (see, for example, [1, §1.13, §1.14]).

From this follows that, if $p, \alpha \in \mathbb{N}$, $\alpha(\mu + 1) - (2p - 1) > 0$, then

$$(4) \quad \sum_{k=1}^{\infty} \frac{2(-1)^{k-1} k^{2p-1}}{(k^{\alpha} + t^{\alpha})^{\mu+1}} \sim -\frac{E_{2p-1}(0)}{t^{\alpha(\mu+1)}} = \frac{(2^{2p} - 1)B_{2p}(0)}{p} \cdot \frac{1}{t^{\alpha(\mu+1)}}, \quad t \rightarrow +\infty.$$

From equality

$$B_{2p}(0) = \frac{(-1)^{p-1}(2p)!}{2^{2p-1}\pi^{2p}} \cdot \sum_{k=1}^{\infty} \frac{1}{k^{2p}}, \quad p \in \mathbb{N},$$

follows that $\frac{(2^{2p}-1)|B_{2p}(0)|}{p} > \frac{(2^{2p}-1)}{p} \cdot \frac{(2p)!}{2^{2p-1}\pi^{2p}} = s_{2p}$, $p \in \mathbb{N}$, where $s_n = \frac{4(2^n-1)}{n^{2n}} \frac{n!}{\pi^n}$, $n \in \mathbb{N}$. From inequality

$$\frac{s_{n+1}}{s_n} = \frac{2^{n+1} - 1}{2^n - 1} \cdot \frac{n}{2\pi} = \left(2 + \frac{1}{2^n - 1}\right) \frac{n}{2\pi} > \frac{n}{\pi} > 1, \quad n \geq 4,$$

follows that $s_n \geq s_8 = \frac{80325}{4\pi^8} = 2,1163 \dots > 2$ for all $n \geq 8$.

Conclusion: if $m, \alpha \in \mathbb{N}$, $\alpha\mu - (4m + 5) > 0$, then inequality

$$\sum_{k=1}^{\infty} \frac{2(-1)^{k-1} k^{4m+5}}{(k^{\alpha} + r^2)^{\mu}} \leq \frac{2}{(1 + r^2)^{\mu}}$$

is not possible for each big enough $r > 0$.

REFERENCES

- [1] H. Bateman, A. Erdélyi, *Higher transcendental functions, vol. 1, 2*, New York, Toronto, London, MC Graw-Hill Book Company, INC, 1953.
- [2] Ž. Tomovski, R. Hilfer, *Some bounds for alternating Mathieu type series*, Journal of Mathematical Inequalities, **2**(1) (2008), p. 17–26.
- [3] V.P. Zastavnyi, *The generalized formula of Euler-Maclaurin and its application*, Transactions of IAMM of NAS of Ukraine, **17** (2008), p. 51–60 (in Russian).
- [4] V.P. Zastavnyi, *Mathieu's series: inequalities, asymptotics and positive definiteness*, <http://arxiv.org/abs/0901.1104v1>, 2009, p. 1–26.

DEPARTMENT OF MATHEMATICS, DONETSK NATIONAL UNIVERSITY, UNIVERSITETSKAYA STR. 24,
340001, DONETSK, UKRAINE

E-mail address: zastavn@rambler.ru